






Legend	
	EMR
	EMT
	Intermediate
	Paramedic
	Medical Control



Poison Control 800-222-1222

<b>EMR</b>	<ol style="list-style-type: none"> <li>1. Consider possible scene &amp; patient contamination and follow agency safety procedures.</li> <li>2. If SAFE to do so, remove from any exposure environment.</li> <li>3. If narcotic overdose suspected, administer <b>Narcan 2-4mg IN</b>. May repeat in 10 min if no response</li> <li>4. Oxygen: Titrate SpO2 to 94-99%. Be prepared to support the patient's respirations with BVM if necessary.</li> <li>5. Gather all medications/pill bottles etc. and give to transport agency.</li> <li>6. Check and record vital signs every 5 minutes, if serious, 15 minutes if minor to moderate, until the transporting unit arrives.</li> </ol>	<b>EMR</b>
------------	--	------------

<b>EMT</b>	<ol style="list-style-type: none"> <li>1. Continue EMR care.</li> <li>2. Apply cardiac monitor and obtain 12-lead EKG if indicated by chief complaint, transmit to receiving facility (if equipped). It is beyond the scope of the EMT to monitor, interpret 12-leads or cardiac rhythms.</li> <li>3. Check Blood glucose level. If &lt; 60, or &lt; 80 with signs/symptoms, treat per Altered Level of Conscious SMO</li> <li>4. Initiate ALS intercept for further medications, if indicated.</li> <li>5. Contact receiving hospital as soon as possible</li> </ol>	<b>EMT</b>
------------	---	------------

<b>I</b>	<ol style="list-style-type: none"> <li>1. Continue EMT care.</li> <li>2. Initiate IV Normal Saline 1-2L bolus to maintain a systolic BP greater than 90 mmHg.</li> <li>3. If BP &lt; 90 mmHg, give 20mL/kg fluid bolus of Normal Saline and re-assess.</li> <li>4. If narcotic overdose suspected, administer <b>Narcan 2-4mg IN?IV/IM</b>. May repeat in 10 min if no response</li> <li>5. Consider ALS intercept should the situation warrant additional medication.</li> <li>6. Contact receiving hospital as soon as possible (need time to prepare) or Medical Control if necessary.</li> </ol>	<b>I</b>
----------	--	----------

<b>P</b>	<ol style="list-style-type: none"> <li>1. Continue EMT-I care.</li> <li>2. If known tricyclic antidepressant (TCA) or Aspirin (ASA) overdose give <b>Sodium Bicarbonate 50mEq IV</b>.</li> <li>3. Transport as soon as possible.</li> <li>4. Contact receiving hospital as soon as possible (need time to prepare) or Medical Control if necessary.</li> </ol>	<b>P</b>
----------	--	----------