

**OB/GYN**

Legend	
	EMR
	EMT
	Intermediate
	Paramedic
	Medical Control

<b>EMR</b>	<ul style="list-style-type: none"> <li>• Determine need for imminent delivery or need for immediate transport;</li> <li>• Position mother for delivery: Have mother lie back, if tolerated, with knees drawn up and spread apart; Elevate buttocks with pillow or blankets;</li> <li>• Whenever possible, use sterile or aseptic technique;</li> <li>• Coach mother to breathe deeply between contractions and to push with contractions;</li> <li>• Administer oxygen using appropriate oxygen delivery device, as indicated per protocol.</li> <li>• Delivery Procedure As the head crowns control with gentle pressure and support the head during delivery; Examine neck for the presence of a looped (nuchal) umbilical cord: If cord is looped around neck, gently slip it over the infant's head. If unable to do so, clamp and cut the cord;</li> </ul>	<b>EMR</b>
<b>EMT</b>	<ul style="list-style-type: none"> <li>• Support the infant's head as it rotates for shoulder presentation: With gentle pressure, guide the infant's head downward to deliver the anterior shoulder and then upward to release the posterior shoulder; Complete the delivery of the infant;</li> <li>• Apply two clamps to umbilical cord (if not already done due to Nuchal cord): The first one is placed approximately ten (10) inches from the infant; The second is placed 2"-3" proximal to the first clamp (7"-8" from infant's abdomen); Cut cord between clamps and check for umbilical cord bleeding;</li> <li>• Umbilical tape may used if umbilical vascular access is being considered (umbilical vein catheterization procedure);</li> <li>• If umbilical cord bleeding is evident apply additional clamp(s) as needed;</li> <li>• Dry infant and wrap in warm towels/blanket (cover infant's head): Place infant on mother's abdomen for mother to hold and support;</li> <li>• Note and record infant's gender, time and geographical location (especially if in transit) of birth: If infant resuscitation is not necessary, record APGAR score at 1 minute and 5 minutes post delivery; If infant resuscitation is necessary, follow neonatal resuscitation protocol; Delivery of the Placenta: (do not delay transport)</li> </ul>	<b>EMT</b>
<b>I</b>	<ul style="list-style-type: none"> <li>• As the placenta delivers, the mother should be encouraged to push with contractions;</li> <li>• Hold placenta with both hands, place in plastic bag or other container and transport with mother to receiving hospital. NEVER "pull on" umbilical cord to assist placenta delivery.</li> </ul>	<b>I</b>

<b>P</b>	<p>Provide additional assessment to all patients (mother and child or multiple children): If newborn presents with signs of distress or hypoperfusion see newborn resuscitation protocol;</p> <p>Establish IV access with sodium chloride (NS) to Keep Vein Open: If patient presents with symptoms/signs of hemodynamic instability, initiate fluid resuscitation per shock protocol with large bore catheter.</p> <p><b>ALS Only:</b> Eclamptic Seizures: Administer <b>midazolam-2-4mg IV/IO or 10mg IM/IN;</b> Consider <b>magnesium sulfate-1-4grams</b> slow IV push over three minutes.</p>	<b>P</b>
----------	--	----------