

Universal Cardiac Care Thinking Points

History:

- Age
- Medications
- Viagra, Levitra, Cialis (within 72hr)
- Past medical history (MI, Angina, Diabetes, Post Menopausal)
- Allergies (Aspirin, Opiates, Lidocaine)
- Recent physical exertion
- Provocation
- Quality (cramp, constant, sharp, dull, etc)
- Radiation
- Severity (1 – 10)
- Time (onset/duration/repetition)
- DNR Status

Signs and Symptoms

- Substernal chest pain/pressure
- Heaviness, tightness or discomfort in chest
- Radiation and/or pain/discomfort to the neck or jaw
- Pain/discomfort/weakness in shoulders/arms
- Consider atypical presentation in diabetics, females, or special populations
- Nausea/vomiting
- Diaphoresis
- Dyspnea
- Time of Onset

Priorities of Care

- Assessing/securing ABCs
- Determine quality/severity of distress
- Identify contributing factors
- Obtaining medical history (including medications & allergies)

Possible Considerations


- Trauma vs. Medical
- Myocardial Infarction
- Pericarditis
- Pulmonary embolism
- Asthma/COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or hiatal hernia
- Esophageal spasm
- Chest wall injury
- Pleural pain
- Tox (Cocaine or Methamphetamine)

Cardiogenic Shock Signs and Symptoms

- Pain heaviness, tightness or discomfort in chest with hypotension (SBP < 100mmHg)
- Rales or crackles (wet lung sounds)
- Pedal edema
- Dyspnea
- Diaphoresis
- Nausea/Vomiting
- Time of Onset


Priorities of Care

- Assessing/securing ABCs
- Determine quality/severity of distress
- Identify contributing factors
- Obtaining medical history (including medications & allergies)



TKO=8-15gtts/min

- V₁ 4th intercostal space to the right of the sternum
- V₂ 4th intercostal space to the left of the sternum directly between the leads V₁ & V₄
- V₃ 5th intercostal space at midclavicular line
- V₄ level with V₄ at left anterior axillary line
- V₅ level with V₄ at left midaxillary line (directly under the midpoint of the amppt)
- V₆ 5th intercostal space, right midclavicular line



Cardiac Arrest Thinking Points

Possible Causes of PEA

- Hypovolemia
- Hypoxia
- Hydrogen ion (acidosis)
- Hypo/Hyperkalemia
- Hypothermia
- Pneumothorax
- Tamponade
- Toxins
- Thrombosis (pulmonary)
- Thrombosis (coronary)

- **If the patient is not conscious and not breathing adequately then start CPR.**
- **If the cardiac arrest is witnessed by EMS personnel, start CPR and defibrillate immediately if indicated.**
- **Compressions should be between 100-120 per minute, ensuring adequate depth and recoil of the chest. There should minimal interruptions in compressions.**
- Patient with implanted pacemakers or defibrillators (AICDs) are treated the same as any other patient.
- A 20mL fluid bolus should be given after each medication to flush the IV line.

- **Treat the patient, not the monitor. A rhythm on the monitor should NOT be used to determine pulse.**
- When changing to ALS monitoring equipment, attach defibrillation cables prior to disconnecting the AED.
- Resuscitation and treatment decisions are based on the duration of the arrest, physical exam and the patient's medical history. Consider cease-effort orders if indicated.
- Consider underlying etiologies and treat according to appropriate protocols.